International Journal of All Research Education and Scientific Methods (IJARESM),
ISSN: 2455-6211, Volume 13, Issue 5, May-2025, Available online at: www.ijaresm.com

The Hidden Burden: Arrhythmias in Hemodialysis Patients and the Pursuit of Cardiovascular Safety

Rose Mary Benny[1], Aldreena Jans[2], Job Sebastian[3], Rosmin Jacob[4]

[1,2,3]Department of Pharmacy, St Jame's College of Pharmaceutical Sciences, Chalakudy, Kerala University of Health and Sciences, India.
[4]Assistant Professor, Department of Pharmacy Practice, St Jame's College of Pharmaceutical Sciences, Chalakudy, Kerala University of Health and Sciences, India

----------**********----------

ABSTRACT

In patients receiving continuous hemodialysis, arrhythmias are a serious but sometimes overlooked complication that significantly increases cardiovascular morbidity and mortality.This cohort has a high incidence of arrhythmic events, with bradyarrhythmia's, including asystole, happening more frequently than previously thought, according to recent research using implantable loop recorders (ILRs). An annualized rate of bradycardia/asystole events, for example, was 0.19 per patient-year, higher than the rate of ventricular tachyarrhythmias, according to a systematic review. A higher risk of sudden cardiac death (SCD) has been linked to factors such low dialysate potassium and calcium concentrations.These results highlight the necessity of increased monitoring and customized treatments to reduce the risk of arrhythmias in hemodialysis patients. Individualized dialysate prescriptions, rigorous electrolyte level monitoring, and enhanced cardiac monitoring in high-risk patients are some possible strategies. To create and validate risk classification instruments and investigate the effectiveness of preventative strategies in this susceptible group, more study is necessary.Particularly during or right after sessions, the timing of these arrhythmias frequently corresponds with the dialysis cycle, indicating that quick changes in electrolyte and fluid balance are crucial.

Keywords: Arrythmia, Cardiovascular Mortality, Chronic Kidney Disease, ESRD, Hemodialysis.

INTRODUCTION

Chronic Kidney Disease (CKD) is a progressive condition marked by a gradual decline in kidney function, which is categorized into five stages based on the severity of the disease. As CKD is increasing rapidly people rely more on hemodialysis as a treatment option for better results.Although hemodialysis is considered a better approach it causes various health concerns to patients of which cardiovascular mortality is a hidden burden [1].Despite advancements in technology this remain unrecognised and poorly controlled and among these arrythmia which is known as abnormal heart rhythm remains a first line cause.

Arrhythmia is now considered a hidden burden among hemodialysis patients. Due to lack of technologies, they are unable to detect effectively,but studies suggest that both atrial and ventricular arrhythmias are reported among these population [2]. As CKD patients are increasing rapidly that depend upon hemodialysis,it is of important concern to reduce this cardiovascular mortality and thereby improve their quality of life.By monitoring each dialysis patients,and thereby analysing their reports,findings indicate that fluctuations in electrolyte levels,fluid overload and various other comorbidities are major cause for cardiovascular disease [3].

EPIDEMIOLOGICAL INSIGHTS INTO DIALYSIS-ASSOCIATED ARRHYTHMIA

About 50 % deaths in hemodialysis occurs due to cardiac arrhythmias which is defined as a group of rhythm disorders and they range from minor, to severe and can lead to serious complications [4]. Among these the most common arrhythmias observed in dialysis patients is atrial fibrillation (AF), which is prevalent approximately in 13%-25% of population.Ventricular arrythmia is also an important arrythmia that are of greater concern due to their association with sudden cardiac arrest [5]. Additionally, bradyarrhythmia's are observed frequently and they lead to complications such as fatigue,syncope etc. The United States Renal Data System (USRDS) and Dialysis Outcomes and Practice Patterns Study (DOPPS) exhibit the findings that prevalence of arrythmias in hemodialysis patients as well as their unrecognised effects [6].

PATHOPHYSIOLOGY: WHY ARE HEMODIALYSIS PATIENTS PRONE TO ARRHYTHMIAS?

The pathophysiology of arrhythmias in hemodialysis patients is complex and linked to both the underlying chronic kidney disease (CKD) and the dialysis procedure itself.

Electrolyte Imbalances

- Potassium: This is one of the important factors causing arrythmia in dialysis patients.During these sessions potassium levels varies and can cause hypokalaemia and hyperkalaemiawhich disrupts normal cardiac functioning.High potassium level can lead to ventricular arrythmias whereas low level led to brady arrhythmia [7].
- Calcium and Magnesium: Fluctuations of calcium and magnesium can lead to arrhythmic episodes. Both low and high levels of calcium can impair cardiac functions and thereby leading to QT interval prolongation and ventricular tachycardia.Magnesium also plays an important role in arrhythmias [8].

Fluid shifts and hemodynamic instability

Hemodialysis can cause fluid shifts and also changes hemodynamics which triggers arrythmias.

- Ultrafiltration is defined as a process of removal of excess fluid from blood during dialysis and there by cause hypovolemia (low blood volume), hypotension, and reduced coronary perfusion which can lead to ischemic arrhythmias [9].
- Fluid overload during dialysis can lead to heart attack and other arrhythmic episodes [10].

Uremic Toxins and Inflammation

Chronic kidney disease can lead to accumulation of toxins in body that can affect heart conducting system.These further lead to:

- Chronic Inflammation which causes fibrosis of myocardium and conducting system.
- Direct myocardial toxicityimpairs cellular functioning and cause arrythmias [11].

RISK FACTORS FOR ARRHYTHMIA IN DIALYSIS PATIENTS

Arrhythmias are observed in hemodialysis patients due to wide range of factors such as clinical conditions, dialysis-related variables, and medication influences.

Structural Heart Disease

Chronic kidney disease (CKD) is a strong factor for structural abnormalities of the heart, such as ischemic heart disease, cardiomyopathy, and valvular disorders. These issues occur due to various contributing factors, such as persistent hypertension, fluid overload, anaemia, and a range of metabolic disturbances related to kidney dysfunction.These structural changes can exacerbate the chances of arrhythmiaandalsoaffects normal conducting system [12].

Medication Effect

Dialysis patients rely on various class of medication for their end stage renal disease as well as the associated comorbidities.Some of the medication that can exacerbate their conditions include Anti-coagulants,Anti-hypertensives and some cardiac drugs like digoxin [13].Even though these medications are essential for managing various disease associated with health they may sometimes worse the current situation directly or indirectly [14].

Dialysis Frequency and Timing

The schedule and frequency of hemodialysis sessions also influence cardiovascular risk as long intradialytic intervals can lead to accumulation of toxins there by lead to cardiac events [15]. The Monday and Tuesday sessions are linked to the highest rates of sudden cardiac death. More Frequent or Daily Dialysis can make more stable electrolyte levels thereby reduce intradialytic hypotension. Nocturnal and Extended Dialysis that is 6-8 hours overnight can reduce hemodynamicstress [16].

TYPES OF ARRHYTHMIAS

There are different types of arrythmia observed in hemodialysis patients due to electrolyte shift and fluid management.Various arrythmias include:

Atrial Fibrillations

- Atrial fibrillation (AF)
  It is the most common sustained arrhythmia in hemodialysis which can be determined through ECG.They can lead to complications such as stroke and other cardiac diseases [17].
- Atrial flutter(AFL)
  They are less common than AF and can be managed by anti-coagulation.

- Atrial Tachycardia(AT)
  It is seen frequently due to volume overload and other electrolyte disturbances [18].

Ventricular Arrhythmias

- Ventricular Tachycardia(VT)
  Mainly occur in patients with underlying heart disease.
- Ventricular Fibrillation(VF)
  This is one of the life-threateningconditions but a less common one which is mainly caused by electrolyte abnormalities such as hyperkalaemia or hypokalaemia [19].

Bradyarrhythmia

- Sinus Bradycardia
  It is defined as sinus rhythm with heart rate greater than 60 beats per minute(bpm).Electrolyte imbalances,medication use, and other autonomic dysfunction can lead to this condition [20].
- Atrioventricular Block(AV)
  Defined as a conduction disorder where electrical signals between atria and ventricles are delayed and commonly observed in hemodialysis patients.Itrequires multidisciplinary approach for the management [21].

CLINICAL MANIFESTATIONS

Asymptomatic
Most of the symptoms of arrythmia in hemodialysis patients remain silent.

Palpitations
Patients experience sudden or rapid heart rhythm which is a common clinical manifestation of arrythmia [22].

Chest Pain or Discomfort
They mainly occur due to reduced coronary hypotension and can exacerbate over time.

Other Symptoms Include:

- Dyspnoea
- Hypotension
- Fatigue or weakness [23]

DIAGNOSTIC CHALLENGES

Diagnostic Tools and Monitoring in Hemodialysis Patients

Electrocardiogram(ECG)
It is one of the diagnostic tools to determine rhythm analysis as well as structural changes of heart.ECGs are mainly conducted during

- Initiation of hemodialysis.
- Routine monitoring in cardiac patients.
- Symptomatic patients should undergo ECG
- After electrolyte correction[24]

Echocardiography
Echocardiography is a non-invasive procedure to determine structural heart changes and cardiovascular mortality.It is mainly performed during post dialysis when the patients is at dry weight.Avoid echoes during pre-dialysis as it may show variations due to fluid overload [25].

Laboratory Tests
Electrolytes and cardiac biomarkers are checked during dialysis inorder to identify fluctuations that may cause arrhythmicevents [26].

Implantable Loop Recorder (ILR)

Long term continuous monitoring device useful in assessing high risk and unexplained syncope cases.ESRD patients have a high risk of arrythmia than general population. Most arrythmias are silent and lethal so difficult to diagnose through ECG monitoring there by this method have an advantage of high sensitivity and can be used for a long term (3-4 hours).Dueto its high cost, they are used in a limited range [27].

Management Strategies And Prevention Of Arrhythmias

Management of arrythmia in hemodialysis patients is essential to reduce cardiovascular mortality and morbidity and thereby improving quality of life.Managing and preventing arrythmia requires multifactorial,multidisciplinary approach [28].Main management includes close monitoring ECG or implantable loop recorders to detect silent,or ischemic arrhythmias.Immediate managementincludes correcting electrolyte imbalances and fluid overload.Long term management includes the use of beta blockers, amiodarone, and anti-coagulants and thereby reducing drugs of high clearance or proarrhythmicdrugs [29].Dialysis specific interventions include avoid rapid ultrafiltration,customise dialysate composition and use a cooler dialysate.Preventive strategies involve controlling risk factorsand thereby reduce sudden cardiac arrest [30].

CONCLUSION

The health and survival of hemodialysis patients are seriously threatened by arrhythmias, which are frequently overlooked. This susceptible group has an alarming rate of sudden cardiac death due to the high prevalence of both symptomatic and silent arrhythmias, which are exacerbated by electrolyte changes, myocardial remodelling, autonomic dysfunction, and stresses associated with dialysis.

Even with improvements in cardiovascular care and dialysis technology, arrhythmic risk is still frequently not adequately identified and mitigated by current treatment approaches. Novel approaches to early detection and tailored intervention include machine learning-based prediction models, biomarker profiling, and continuous cardiac monitoring. A more proactive and preventive approach also requires reassessing dialysis procedures, improving medication, and encouraging interdisciplinary cooperation between cardiologists and nephrologists. Future research must prioritize not only understanding the mechanistic underpinnings of arrhythmogenesis in end-stage renal disease but also translating these insights into practical, scalable solutions. By bringing arrhythmias to the forefront of dialysis-related care and embracing innovations in monitoring and risk management, we can begin to address this hidden burden and move closer to ensuring cardiovascular safety for all patients on hemodialysis.

REFERENCES

[1]. Firas Ajam, Swapnil Patel,Anas Alrefaee, DawnCalderon,Mohammad Hossain,Arif AsifCardiac Arrhythmias in Patients with End Stage Renal Disease (ESRD) on Hemodialysis American Journal of Internal Medicine. Vol. 7, No. 1, 2019, pp. 22-26

[2]. Philip A Kalra, Darren Green, Dimitrios Poulikakos Arrhythmia in hemodialysis patients and its relation to sudden death Kidney international 93 (4), 781-783, 2018

[3]. Luminita Voroneanu, Adrian Covic Arrhythmias in hemodialysis patients JN journal of nephrology 11 (6), 716, 2009

[4]. Simonetta Genovesi, Antonio Vincenti, Emanuela Rossi, Daniela Pogliani, Irene Acquistapace, Andrea Stella, Maria Grazia Valsecch Atrial fibrillation and morbidity and mortality in a cohort of long-term hemodialysis patients American Journal of Kidney Diseases 51 (2), 255-262, 2008

[5]. Ken-ichi Kimura, Kaoru Tabei, Yasushi Asano, Saichi Hosoda Cardiac arrhythmias in hemodialysis patients: A study of incidence and contributory factors Nephron 53 (3), 201-207, 1989

[6]. Volker Wizemann, Lin Tong, Sudtida Satayathum, Alex Disney, Takashi Akiba, Rachel B Fissell, Peter G Kerr, Eric W Young, Bruce M Robinson Atrial fibrillation in hemodialysis patients: clinical features and associations with anticoagulant therapy Kidney international 77 (12), 1098-1106, 2010

[7]. Paul R Roberts, Kurt Stromberg, Lawrence C Johnson, Benedict M Wiles, Thomas A Mavrakanas, David M Charytan A systematic review of the incidence of arrhythmias in hemodialysis patients undergoing long-term monitoring with implantable loop recorders Kidney International Reports 6 (1), 56-65, 2021

[8]. Gail Morrison, Eric L Michelson, Steven Brown, Joel Morganroth Mechanism and prevention of cardiac arrhythmias in chronic hemodialysis patients Kidney international 17 (6), 811-819, 1980

[9]. Firas Ajam, Swapnil Patel, Anas Alrefaee, Dawn Calderon, Mohammad A Hossain, Arif Asi Cardiac Arrhythmias in Patients with End Stage Renal Disease (ESRD) on Hemodialysis; Recent Update and Brief American Journal of Internal Medicine 7 (1), 22-26, 2019

[10]. Wolfgang C Winkelmayer, Amanda R Patrick, Jun Liu, M Alan Brookhart, Soko Setoguchi Wolfgang C Winkelmayer, Amanda R Patrick, Jun Liu, M Alan Brookhart, Soko Setoguchi Journal of the American Society of Nephrology 22 (2), 349-357, 2011

[11]. Oz M Shapira, Yaacov Bar-Khayim ECG changes and cardiac arrhythmias in chronic renal failure patients on hemodialysis Journal of electrocardiology 25 (4), 273-279, 1992
[12]. P Meier, P Vogt, E Blan Ventricular arrhythmias and sudden cardiac death in end-stage renal disease patients on chronic hemodialysis Nephron 87 (3), 199-214, 2001
[13]. A Blumberg, M Häusermann, B Strub, HR Jenzer Cardiac arrhythmias in patients on maintenance hemodialysis Nephron 33 (2), 91-95, 1983
[14]. Frederic Sacher, Laurence Jesel, Claire Borni-Duval, Valerie De Precigout, Frédéric Lavainne, Jean-Philippe Bourdenx et.al Cardiac rhythm disturbances in hemodialysis patients: early detection using an implantable loop recorder and correlation with biological and dialysis parameters JACC: Clinical Electrophysiology 4 (3), 397-408, 2018
[15]. Eleni Stamellou, Christos Georgopoulos, Lampros Lakkas, Evangelia Dounous 10 tips on how to manage severe arrhythmia in hemodialysis patients Clinical Kidney Journal, sfaf072, 2025
[16]. An S De Vriese, Gunnar Heine Anticoagulation management in haemodialysis patients with atrial fibrillation: evidence and opinion Nephrology Dialysis Transplantation 37 (11), 2072-2079, 2022
[17]. Rahul Samanta, Christopher Chan, Vijay S Chauhan Arrhythmias and sudden cardiac death in end stage renal disease: epidemiology, risk factors, and management Canadian Journal of Cardiology 35 (9), 1228-1240, 2019
[18]. Nichole M Rogovoy, Stacey J Howell, Tiffany L Lee, Christopher Hamilton, Erick A Perez-Alday, Muammar M Kabir et.al Hemodialysis Procedure–Associated Autonomic Imbalance and Cardiac Arrhythmias: Insights from Continuous 14-Day ECG Monitoring Journal of the American Heart Association 8 (19), e013748, 2019
[19]. Mehmet Kanbay, Baris Afsar, David Goldsmith et.al Sudden death in hemodialysis: an update Blood purification 30 (2), 135-145, 2010
[20]. Paul R Roberts, Darren Green Arrhythmias in chronic kidney disease Heart 97 (9), 766-773, 2011
[21]. Huseyin Bozbas, Ilyas Atar, Aylin Yildirir, AliseydiOzgul, Murathan Uyar, Nurhan Ozdemir, Haldun Muderrisoglu, Bulent Ozin Prevalence and predictors of arrhythmia in end stage renal disease patients on hemodialysis Renal failure 29 (3), 331-339, 2007
[22]. Prabir Roy-Chaudhury, Jim A Tumlin, Bruce A Koplan et.al Primary outcomes of the Monitoring in Dialysis Study indicate that clinically significant arrhythmias are common in hemodialysis patients and related to dialytic cycle Kidney international 93 (4), 941-951, 2018
[23]. Ajit S Narula, Vivekanand Jha, Harinder K Bali, Vinay Sakhuja Cardiac arrhythmias and silent myocardial ischemia during hemodialysis Renal failure 22 (3), 355-368, 2000
[24]. James O Burton, ShvanKorsheed, Ben J Grundy, Christopher W McIntyre Hemodialysis-induced left ventricular dysfunction is associated with an increase in ventricular arrhythmias Renal failure 30 (7), 701-709, 2008
[25]. Mansur Kayatas, Arif Ertürk, Haldun Muderrisoglu, Emin Korkmaz, Nurhan Özdemir Acetate hemodialysis does not increase the frequency of arrhythmia in hemodialysispatient's Artificial organs 22 (9), 781-784, 1998
[26]. Hamidreza Nasri, Behzad Sarvar Azimzadeh, MohamadhoseinTorabinejad, Reza Pourakbari Cardiac arrhythmia in dialysis patients ARYA Atherosclerosis Journal 3 (4), 2010
[27]. Simonetta Genovesi, Antonio Vincenti, Emanuela Rossi, Daniela Pogliani, Irene Acquistapace, Andrea Stella et.al Atrial fibrillation and morbidity and mortality in a cohort of long-term hemodialysis patients American Journal of Kidney Diseases 51 (2), 255-262, 2008
[28]. Yamita Sakhare, Alan Almeida, Deepak Phalgune, Aditi Erande, Sanjay Madhav Mehendale The Frequency, Causes and Patterns of Asymptomatic Cardiac Arrhythmias in Patients on Maintenance Hemodialysis Indian Journal of Nephrology 35 (3), 397, 2025
[29]. James A Tumlin, Prabir Roy-Chaudhury, Bruce A Koplan et.al Relationship between dialytic parameters and reviewer confirmed arrhythmias in hemodialysis patients in the monitoring in dialysis study BMC nephrology 20, 1-13, 2019
[30]. Antonio Santoro, Elena Mancini, Gerad London, Lucile Mercadal, Hafedh Fessy, Bruno Perrone, Leonardo Cagnoli, Eleonora Grandi et.al Patients with complex arrhythmias during and after haemodialysis suffer from different regimens of potassium removal Nephrology Dialysis Transplantation 23 (4), 1415-1421, 2008